

SOUTH DAKOTA STATE UNIVERSITY College of Agriculture, Food and Environmental Sciences

South Dakota State University Extension South Dakota Agricultural Experiment Station at SDSU

2023 South Dakota Conventional Soybean Variety Trial Results Brookings

Jonathan KleinjanSDSU Extension AgronomistKevin KirbyAgricultural Research ManagerShawn HawksAgricultural Research Manager

Location:	3 miles north of Brookings (57006) in Brookings County, SD 44.362528°, -96.792450°				
Cooperator:	SDSU Felt Family Research Farm				
Soil Type:	Vienna-Brookings complex, 0-2% slopes				
Fertilizer:	none				
Previous crop:	corn				
Tillage:	conventional				
Row spacing:	30 inches				
Seeding Rate:	150,000/acre				
Herbicide:	Pre: 1 pt/acre Dual II Magnum Post: 12 oz/acre Volunteer + 1 pt/acre Ultra Blazer				
Insecticide:	none				
Date seeded:	5/17/2023				
Date harvested:	10/17/2023				

SDSU Extension is an equal opportunity provider and employer in accordance with the nondiscrimination policies of South Dakota State University, the South Dakota Board of Regents and the United States Department of Agriculture.



2023 South Dakota Conventional Soybean Variety Trial Results Brookings

SOUTH DAKOTA STATE UNIVERSITY EXTENSION

Table 1. Conventional soybean performance results (average of 4 replications - Maturity Groups 0 & I) at Brookings, SD.

Variety Information		Agronomic Performance			
Brand	Variety	Maturity Rating	Yield (bu/ac@13%)	Moisture (%)	Lodging Score (1-5)*
Viking Seed	VK 0821N	0.8	76.9	10.6	2.3
Ag Performance	AP 0720	1.0	75.8	10.5	2.0
Viking Seed	VK 1223N	1.2	72.3	10.3	1.3
Miller Hybrids	1938	1.9	67.8	10.1	2.8
Viking Seed	VK 1518N	1.5	67.4	10.4	1.0
LG Seeds	LGS1684	1.6	67.2	10.4	1.5
Check	AG15XF2	1.5	65.6	10.5	2.5
Miller Hybrids	1538	1.5	64.4	10.4	1.8
Brushvale Seed	BS1799	1.5	64.1	10.3	1.5
Ag Performance	AP 1321	1.3	63.9	10.6	2.0
SD AES	BROOKINGS	1.7	63.3	10.7	2.5
Ag Performance	AP 1621	1.6	63.2	10.5	2.3
Brushvale Seed	BS91614	1.0	63.0	10.3	1.0
Virtue	V1621	1.6	62.8	10.0	1.0
Brushvale Seed	BS1940	1.9	62.2	10.3	1.5
Viking Seed	VK 1718N	1.7	60.6	10.3	1.0
Richland IFC	MK146	1.1	60.0	10.3	1.3
Richland IFC	MK41	1.1	59.6	10.5	3.5
Richland IFC	MK0603	0.6	57.6	10.7	3.5
Richland IFC	MK808CN	0.8	57.2	11.0	3.3
MN AES	MN1807CN	1.8	56.1	10.7	2.0
Richland IFC	MK1023	1.0	55.7	10.5	1.8
Richland IFC	MK1423	1.4	52.5	10.4	1.8
Richland IFC	MK9103	1.3	44.5	10.2	2.3
Richland IFC	SABLE	1.2	43.8	10.1	3.8
Richland IFC	DECKER	1.0	43.3	9.7	1.5
Richland IFC	MK9102	1.2	42.9	10.5	3.0
Richland IFC	MK9101	1.1	41.8	10.9	2.5
		Trial Average	59.9	10.4	2.0
LSD (0.05)†			3.2	0.2	-
		C.V.‡	3.8	-	-

* Lodging Score (1 = no lodging to 5 = flat on the ground)

+ Yield or moisture value required (>LSD) to determine if varieties are significantly different from one another.

+ C.V. is a measure of variability or experimental error, 15% or less is acceptable.



2023 South Dakota Conventional Soybean Variety Trial Results Brookings

SOUTH DAKOTA STATE UNIVERSITY EXTENSION

Table 2. Conventional soybean performance results (average of 4 replications - Maturity Group II) at Brookings, SD.

Variety Information		Agronomic Performance				
Brand	Variety	Maturity Rating	Yield (bu/ac@13%)	Moisture (%)	Lodging Score (1-5)*	
Check	AG15KF2	1.5	63.6	10.4	1.0	
Viking Seed	VK 2022N	2.0	60.4	10.5	1.0	
LG Seeds	LGS2020	2.0	59.8	10.2	1.0	
Virtue	V2122	2.1	58.4	10.6	1.0	
Miller Hybrids	2238	2.2	56.6	10.4	1.0	
Virtue	V2922	2.9	55.2	10.3	1.0	
LG Seeds	LGS2329	2.3	53.9	11.0	1.0	
Richland IFC	MK373	2.0	53.2	10.7	1.0	
Viking Seed	VK 2155N	2.1	51.2	10.6	1.0	
Trial Average			56.9	10.5	1.0	
LSD (0.05)†			2.0	0.6	-	
C.V.‡			2.4	-	-	

* Lodging Score (1 = no lodging to 5 = flat on the ground)

+ Yield or moisture value required (≥LSD) to determine if varieties are significantly different from one another.

+ C.V. is a measure of variability or experimental error, 15% or less is acceptable.